

# Hepatitis C Virus Infection, Past or Present

## 2005 Case Definition

### Clinical description

Most HCV-infected persons are asymptomatic. However, many have chronic liver disease, which can range from mild to severe including cirrhosis and liver cancer.

### Laboratory criteria for diagnosis

- Anti-HCV positive (repeat reactive) by EIA, verified by an additional more specific assay (e.g. RIBA for anti-HCV or nucleic acid testing for HCV RNA),

OR

- HCV RIBA positive,

OR

- Nucleic acid test for HCV RNA positive,

OR

- Report of HCV genotype

OR

- Anti-HCV screening-test-positive with a signal to cut-off ratio predictive of a true positive as determined for the particular assay (e.g.,  $\geq 3.8$  for the enzyme immunoassays) as determined and posted by CDC.

### Case classification

*Probable:* a case that is anti-HCV positive (repeat reactive) by EIA and has alanine aminotransferase (ALT or SGPT) values above the upper limit of normal, but the anti-HCV EIA result has not been verified by an additional more specific assay or the signal to cutoff ratio is unknown.

*Confirmed:* a case that is laboratory confirmed and that does not meet the case definition for acute hepatitis C.

**See also:**

- 2003 Case Definition [Hepatitis C Virus Infection (Past or Present)]